

Nuclear and Particle Physics

Dr. Dan Protopopescu

Kelvin Building, room 524

Dan.Protopopescu@glasgow.ac.uk

Recap puzzle


Concept	Equation	Definition
Exponential decay	$N(t) = N_0 e^{-\lambda t}$	Number of nuclei that have decayed in the time t
Activity	$A(t) = A_0 e^{-\lambda t}$	Number of nuclei decaying per unit time
Decay probability	$P_{\text{decay}}(t) = \lambda e^{-\lambda t}$	Probability of a single nucleus decaying in the interval t
Lifetime	$\tau = 1/\lambda$	Maximum time until an unstable nucleus decays
Half-life	$t_{1/2} = \ln 2/\lambda$	Time by which half the radioactive sample has not yet decayed

Answers

Concept	Equation	Definition
Exponential decay	$N(t) = N_0 e^{-\lambda t}$	Number of nuclei that have not decayed by time t
Activity	$A(t) = A_0 e^{-\lambda t}$	Number of nuclei decaying per unit time, where $A_0 = \lambda N_0$
Decay probability	$P_{\text{decay}}(t) = \lambda e^{-\lambda t}$	Probability of a single nucleus decaying in the interval $t \rightarrow t+dt$
Mean lifetime or simply <i>lifetime</i>	$\tau = 1/\lambda$	Mean time until an unstable nucleus decays
Half-life	$t_{1/2} = \ln 2 / \lambda$	Time after which half the radioactive sample has decayed

Simple decay

If a sample of material consists of nucleus A which is unstable and decays to nucleus B (of which there are initially none) we have simply:



Nomenclature:

A - “parent”

B - “daughter”

The initial number of each nucleus is:

$$N_A(t = 0) = N_0 \quad (= \text{total number of nuclei})$$

$$N_B(t = 0) = 0$$

As nucleus A decays into nucleus B

$$N_A(t) = N_0 e^{-\lambda_A t}$$

and since

$$N_0 = N_A(t) + N_B(t)$$

$$N_B(t) = N_0 (1 - e^{-\lambda_A t})$$

Alternative decay modes

An initial nuclide A that decays into two products: $A \rightarrow B + C$

We have at any time t : $N_A(t) + N_B(t) + N_C(t) = N_0$ and

$$\frac{dN_A}{dt} = -\lambda_A N_A, \quad \frac{dN_B}{dt} = \lambda_B N_A, \quad \frac{dN_C}{dt} = \lambda_C N_A$$

with $\lambda_A = \lambda_B + \lambda_C$. The decay constants λ_B and λ_C only determine the probabilities of the decays to products B or C

$$N_B(t) = \frac{\lambda_B}{\lambda_A} N_0 (1 - e^{-\lambda_A t})$$

$$N_C(t) = \frac{\lambda_C}{\lambda_A} N_0 (1 - e^{-\lambda_A t})$$

and

$$N_A(t) = N_0 - N_B(t) - N_C(t) = N_0 e^{-\lambda_A t}$$

Decay series (or chains)

Many heavy nuclei decay via complicated series involving several α and β decays. Consider the simple case of $A \rightarrow B \rightarrow C$, where C is stable and only A is present initially:

The number of nuclei A vary according to:

$$N_A(t) = N_0 e^{-\lambda_A t}$$

The number of nuclei B as a function of time can be found from:

$$\frac{dN_B(t)}{dt} = -\lambda_B N_B(t) + \lambda_A N_A(t) \quad (1)$$



where the first term is the decay of nuclei B and the second term is due to B being created from the decay of A.

Integrating, we can get $N_B(t)$ and its activity $A_B(t)$:

$$N_B(t) = \frac{\lambda_A}{\lambda_B - \lambda_A} N_0 (e^{-\lambda_A t} - e^{-\lambda_B t}) \quad (2)$$

$$A_B(t) = \lambda_B N_B(t) = \frac{\lambda_A \lambda_B}{\lambda_B - \lambda_A} N_0 (e^{-\lambda_A t} - e^{-\lambda_B t})$$

How was equation (2) derived ?

We multiply both sides of the equation by $e^{\lambda_B t}$

$$\frac{dN_B(t)}{dt} = -\lambda_B N_B(t) + \lambda_A N_A \quad | \times e^{\lambda_B t}$$

and we rearrange to obtain

$$e^{\lambda_B t} \frac{dN_B(t)}{dt} + \lambda_B e^{\lambda_B t} N_B(t) = \lambda_A N_A e^{\lambda_B t}$$

This can be written as

$$\frac{d}{dt} (e^{\lambda_B t} N_B(t)) = \lambda_A N_A e^{\lambda_B t}$$

where we use $N_A(t) = N_0 e^{-\lambda_A t}$ to obtain the form

$$\frac{d}{dt} (e^{\lambda_B t} N_B(t)) = \lambda_A N_0 e^{(\lambda_B - \lambda_A)t}$$

How was eq.(2) derived (part II)

We multiply by dt and integrate both sides

$$\int_0^t d(e^{\lambda_B t} N_B(t)) = \int_0^t \lambda_A N_0 e^{(\lambda_B - \lambda_A)t} dt$$

to obtain

$$N_B(t)e^{\lambda_B t} - 0 = \frac{\lambda_A}{\lambda_B - \lambda_A} N_0 (e^{(\lambda_B - \lambda_A)t} - 1)$$

which gives us

$$N_B(t) = \frac{\lambda_A}{\lambda_B - \lambda_A} N_0 (e^{-\lambda_A t} - e^{-\lambda_B t})$$

QED

A → B → C decay series

For the stable element C from such a series one would obtain:

$$N_C(t) = N_0 \left[1 - \frac{\lambda_B e^{-\lambda_A t} - \lambda_A e^{-\lambda_B t}}{\lambda_B - \lambda_A} \right]$$

which we derived using $N_0 = N_A(t) + N_B(t) + N_C(t)$

Instead we will focus on $N_B(t)$ and investigate a few special cases:

- $\lambda_A \gg \lambda_B$ (parent decays quickly)
- $\lambda_A = \lambda_B$
- $\lambda_A < \lambda_B$
- $\lambda_A \ll \lambda_B$ (parent is long lived)

A → B → C decay series for $\lambda_A \gg \lambda_B$

Parent decays quickly, $\tau_A \ll \tau_B$

The number of daughter nuclei rises to maximum, then decays with constant λ_B .


$$N_B(t) = \frac{\lambda_A}{\lambda_B - \lambda_A} N_0 (e^{-\lambda_A t} - e^{-\lambda_B t}) \xrightarrow{\lambda_A \gg \lambda_B} N_0 e^{-\lambda_B t}$$

Annotations in the image: $\lambda_A \gg \lambda_B$ above the fraction; $=-1$ below the denominator; $=0$ below the $e^{-\lambda_A t}$ term.

After a given time, daughter nuclei decay almost as if there were no parent nuclei.

A → B → C decay series for $\lambda_A = \lambda_B$

The solution of eq.(1) when $\lambda_A = \lambda_B = \lambda$ is: $N_B(t) = \lambda N_0 t e^{-\lambda t}$



A → B → C decay series for $\lambda_A < \lambda_B$

Ratio of activities becomes constant after sufficiently long time:

$$\begin{aligned}\frac{A_B}{A_A} &= \frac{\lambda_B N_B}{\lambda_A N_A} = \frac{\lambda_B}{\lambda_B - \lambda_A} (1 - e^{-(\lambda_B - \lambda_A)t}) \\ &\approx \frac{\lambda_B}{\lambda_B - \lambda_A} \quad \text{when } t \rightarrow \infty\end{aligned}$$

A → B → C decay series for $\lambda_A \ll \lambda_B$

Parent nucleus is long lived: $\lambda_A \ll \lambda_B$ or $\tau_A \gg \tau_B$ so:

$$e^{-\lambda_A t} \approx 1 \Rightarrow N_A \approx N_0$$

$$\Rightarrow N_B \approx N_A \frac{\lambda_A}{\lambda_B} (1 - e^{-\lambda_B t})$$

After a sufficiently long time $(1 - e^{-\lambda_B t}) \rightarrow 1$

$$\Rightarrow \lambda_A N_A = \lambda_B N_B \iff dN_B / dt = 0 \quad \text{in eq. (1)}$$

Activity of A = Activity of B

This is known as a *secular equilibrium*, i.e. at large times B is decaying at the same rate as it is produced.

Secular equilibrium ($\lambda_A \ll \lambda_B$)

An example of secular equilibrium is:





Figure 6.6 In the decay ^{132}Te (78 h) \rightarrow ^{132}I (2.28 h) \rightarrow ^{132}Xe , approximate secular equilibrium is reached at about 12 h.

Alpha decay chains

Because α -decay always decreases the atomic mass number A of the nucleus by 4, almost any decay will result in a nucleus with an atomic mass A' such that

$$A \bmod 4 = A' \bmod 4$$

As a result, there are four radioactive decay chains known as the **Thorium** ($4n$), **Neptunium** ($4n+1$), **Radium** ($4n+2$) and **Actinium** ($4n+3$) series.





Image credits: Wikipedia

Thorium series and the age of the Earth



^{232}Th has a very long half life ($t_{1/2} = 14\text{Gyr}$) and goes through a long decay chain to stable ^{208}Pb .

It effectively behaves as if $^{232}\text{Th} \rightarrow ^{208}\text{Pb}$

By measuring the relative abundance of ^{208}Pb :

$$\frac{N(^{208}\text{Pb})}{N(^{232}\text{Th})} = \frac{N_0(1 - e^{-\lambda_{\text{Th}}t})}{N_0 e^{-\lambda_{\text{Th}}t}}$$

one can estimate of the age of the Earth at $4 \times 10^9 \text{yr}$.

Image credits: Wikipedia

Radiometric dating

- Technique used to date geological materials (rocks) or man-made materials
- Based on a comparison between the observed abundance of a naturally occurring radioactive isotope and its decay products, using known decay rates.

<i>Isotope</i>		<i>Half Life (years)</i>	<i>Useful Range (years)</i>	<i>Rock or mineral host</i>
<i>Parent</i>	<i>Daughter</i>			
U-238	Pb-206	4.5×10^9	$10^7 - 4.6 \times 10^9$	Zircon Uraninite
K-40	A-40	1.3×10^9	$5 \times 10^4 - 4.6 \times 10^9$	Micas Hornblende Volcanics
Rb-87	Sr-87	47×10^9	$10^7 - 4.6 \times 10^9$	Micas Orthoclase Igneous rocks
C-14	(N-14)	5730	$100 - 7 \times 10^4$	Biol material CO ₂

Radiocarbon dating

- Carbon is a fundamental part of living tissue.
- There are 3 isotopes of carbon - ^{12}C , ^{13}C and ^{14}C - in the atmosphere, from where they are absorbed by living organisms.
 - The ratio of $^{14}\text{C}/^{12}\text{C}$ is known to be $\gamma_0 = 1.8 \times 10^{-12}$
 - ^{14}C is permanently created by cosmic rays, i.e. this isotopic ratio is constant in nature
- The concentration of ^{14}C in living organisms is the same as that in the environment
- When the organism dies it no longer absorbs ^{14}C . The ^{14}C in the organism decays but the amount of ^{12}C remains constant

$$^{14}\text{C}/^{12}\text{C} = \gamma = \gamma_0 e^{-\lambda t}$$

- By measuring the ratio of $^{14}\text{C}/^{12}\text{C}$ one can find out how much time has passed

$$t = \lambda \ln(\gamma_0/\gamma)$$

